

Genetically Modified Animals in Endocrinology

The following is the latest addition to the Genetically Modified Animals in Endocrinology Bibliography Project. To see earlier segments of this project and links to other useful sources, go to <http://www.endo-society.org/journals/bibliographies.cfm>

Please send your comments, suggestions, and offerings of other relevant citations to this e-mail address: ERKO@endo-society.org

Knockout Mouse Models for Bone Studies*

Bone Morphogenetic Proteins

Dunn NR, Winnier GE, Hargett LK, Schrick JJ, Fogo AB, Hogan BL 1997 Haploinsufficient phenotypes in Bmp4 heterozygous null mice and modification by mutations in Gli3 and Alx4. *Dev Biol* 88:235–247

Bailon-Plaza A, Lee AO, Veson EC, Farnum CE, van der Meulen MC 1999 BMP-5 deficiency alters chondrocytic activity in the mouse proximal tibial growth plate. *Bone* 24:211–216

Jena N, Martin-Seisdedos C, McCue P, Croce CM 1997 BMP7 null mutation in mice: developmental defects in skeleton, kidney, and eye. *Exp Cell Res* 230:28–37

Dudley AT, Robertson EJ 1997 Overlapping expression domains of bone morphogenetic protein family members potentially account for limited tissue defects in BMP7 deficient embryos. *Dev Dyn* 208:349–362

Smad Family

Borton AJ, Frederick JP, Datto MB, Wang XF, Weinstein RS 2001 The loss of Smad3 results in a lower rate of bone formation and osteopenia through dysregulation of osteoblast differentiation and apoptosis. *J Bone Miner Res* 16:1754–1764

Chang H, Huylebroeck D, Verschueren K, Guo Q, Matzuk MM, Zwijsen A 1999 Smad5 knockout mice die at mid-gestation due to multiple embryonic and extraembryonic defects. *Development* 126:1631–1642

TGF β Family

Hamrick MW, McPherron AC, Lovejoy CO 2002 Bone mineral content and density in the humerus of adult myostatin-deficient mice. *Calcif Tissue Int* 71:63–68

Calcitonin

Hoff AO, Catala-Lehnen P, Thomas PM, Priemel M, Rueger JM, Nasonkin I, Bradley A, Hughes MR, Ordonez N, Cote GJ, Amling M, Gagel RF 2002 Increased bone mass is an unexpected phenotype associated with deletion of the calcitonin gene. *J Clin Invest* 110:1849–1857

PTH

Miao D, He B, Karaplis AC, Goltzman D 2002 Parathyroid hormone is essential for normal fetal bone formation. *J Clin Invest* 109:1173–1182

Kitahara Y, Suda N, Kuroda T, Beck F, Hammond VE, Takano Y 2002 Disturbed tooth development in parathyroid hormone-related protein (PTHrP)-gene knockout mice. *Bone* 30:48–56

Suda N, Baba O, Udagawa N, Terashima T, Kitahara Y, Takano Y, Kuroda T, Senior PV, Beck F, Hammond VE 2001 Parathyroid hormone-related protein is required for normal intramembranous bone development. *J Bone Miner Res* 16:2182–2191

Lanske B, Amling M, Neff L, Guiducci J, Baron R, Kronenberg HM 1999 Ablation of the PTHrP gene or the PTH/PTHrP receptor gene leads to distinct abnormalities in bone development. *J Clin Invest* 104:399–407

TNF Family/Osteoprotegerin

Mizuno A, Amizuka N, Irie K, Murakami A, Fujise N, Kanno T, Sato Y, Nakagawa N, Yasuda H, Mochizuki S, Gomibuchi T, Yano K, Shima N, Washida N, Tsuda E, Morinaga T, Higashio K, Ozawa H 1998 Severe osteoporosis in mice lacking osteoclastogenesis inhibitory factor/osteoprotegerin. *Biochem Biophys Res Commun* 247:610–615

Bucay N, Sarosi I, Dunstan CR, Morony S, Tarpley J, Capparelli C, Scully S, Tan HL, Xu W, Lacey DL, Boyle WJ, Simonet WS 1998 Osteoprotegerin-deficient mice develop early onset osteoporosis and arterial calcification. *Genes Dev* 12:1260–1268

Fata JE, Kong YY, Li J, Sasaki T, Irie-Sasaki J, Moorehead RA, Elliott R, Scully S, Voura EB, Lacey DL, Boyle WJ, Khokha R, Penninger JM 2000 The osteoclast differentiation factor osteoprotegerin-ligand is essential for mammary gland development. *Cell* 103:41–50

Lomaga MA, Yeh WC, Sarosi I, Duncan GS, Furlonger C, Ho A, Morony S, Capparelli C, Van G, Kaufman S, van der Heiden A, Itie A, Wakeham A, Khoo W, Sasaki T, Cao Z, Penninger JM, Paige CJ, Lacey DL, Dunstan CR, Boyle WJ, Goeddel DV, Mak TW 1999 TRAF6 deficiency results in osteopetrosis and defective in-

* We thank Rheem D. Medh for compiling this bibliography. Dr. Medh is with the Department of Biology at California State University, Northridge.

terleukin-1, CD40, and LPS signaling. *Genes Dev* 13:1015–1024

Naito A, Azuma S, Tanaka S, Miyazaki T, Takaki S, Takatsu K, Nakao K, Nakamura K, Katsuki M, Yamamoto T, Inoue J 1999 Severe osteopetrosis, defective interleukin-1 signalling and lymph node organogenesis in TRAF6-deficient mice. *Genes Cells* 4:353–362

Yamashita T, Okada S, Higashio K, Nabeshima Y, Noda M 2002 Double mutations in *klotho* and osteoprotegerin gene loci rescued osteopetrotic phenotype. *Endocrinology* 143:4711–4717

Osteopontin

Liaw L, Birk DE, Ballas CB, Whitsitt JS, Davidson JM, Hogan BL 1998 Altered wound healing in mice lacking a functional osteopontin gene (*spp1*). *J Clin Invest* 101:1468–1478

Boskey AL, Spevak L, Paschalis E, Doty SB, McKee MD 2002 Osteopontin deficiency increases mineral content and mineral crystallinity in mouse bone. *Calcif Tissue Int* 71:145–514 (Epub 2002 Jun 20)

Kitahara K, Ishijima M, Rittling SR, Tsuji K, Kurosawa H, Nifuji A, Denhardt DT, Noda M 2003 Osteopontin deficiency induces parathyroid hormone enhancement of cortical bone formation. *Endocrinology* 144:2132–2140

Yoshitake H, Rittling SR, Denhardt DT, Noda M 1999 Osteopontin-deficient mice are resistant to ovariectomy-induced bone resorption. *Proc Natl Acad Sci USA* [Erratum (1999) 96:10944] 96:8156–8160

NF- κ B

Franzoso G, Carlson L, Xing L, Poljak L, Shores EW, Brown KD, Leonardi A, Tran T, Boyce BF, Siebenlist U 1997 Requirement for NF- κ B in osteoclast and B-cell development. *Genes Dev* 11:3482–3496

Iotsova V, Caamano J, Loy J, Yang Y, Lewin A, Bravo R 1997 Osteopetrosis in mice lacking NF- κ B1 and NF- κ B2. *Nat Med* 3:1285–1289

Xing L, Carlson L, Story B, Tai Z, Keng P, Siebenlist U, Boyce BF 2003 Expression of either NF- κ B p50 or p52 in osteoclast precursors is required for IL-1-induced bone resorption. *J Bone Miner Res* 18:260–269

Estrogen Receptor/Androgen Receptor

Sims NA, Dupont S, Krust A, Clement-Lacroix P, Minet D, Resche-Rigon M, Gaillard-Kelly M, Baron R 2002 Deletion of estrogen receptors reveals a regulatory role for estrogen receptors- β in bone remodeling in females but not in males. *Bone* 30:18–25

Windahl SH, Hollberg K, Vidal O, Gustafsson JA, Ohlsson C, Andersson G 2001 Female estrogen receptor β -/- mice are partially protected against age-related trabecular bone loss. *J Bone Miner Res* 16:1388–1398

Lindberg MK, Alatalo SL, Halleen JM, Mohan S, Gustafsson JA, Ohlsson C 2001 Estrogen receptor spec-

ificity in the regulation of the skeleton in female mice. *J Endocrinol* 171: 229–236

Yeh S, Tsai MY, Xu Q, Mu XM, Lardy H, Huang KE, Lin H, Yeh SD, Altuwajri S, Zhou X, Xing L, Boyce BF, Hung MC, Zhang S, Gan L, Chang C, Hung MC 2002 Generation and characterization of androgen receptor knock-out (ARKO) mice: an in vivo model for the study of androgen functions in selective tissues. *Proc Natl Acad Sci USA* [Erratum (2002) 99:15245] 99:13498–13503 (Epub 2002 Oct 07)

Vitamin D

Yoshizawa T, Handa Y, Uematsu Y, Takeda S, Sekine K, Yoshihara Y, Kawakami T, Arioka K, Sato H, Uchiyama Y, Masushige S, Fukamizu A, Matsumoto T, Kato S 1997 Mice lacking the vitamin D receptor exhibit impaired bone formation, uterine hypoplasia and growth retardation after weaning. *Nat Genet* 16:391–396

Yagishita N, Yamamoto Y, Yoshizawa T, Sekine K, Uematsu Y, Murayama H, Nagai Y, Krezel W, Chambon P, Matsumoto T, Kato S 2001 Aberrant growth plate development in VDR/RXR γ double null mutant mice. *Endocrinology* 142:5332–5341

Safadi FF, Thornton P, Magiera H, Hollis BW, Gentile M, Haddad JG, Liebhaber SA, Cooke NE 1999 Osteopathy and resistance to vitamin D toxicity in mice null for vitamin D binding protein. *J Clin Invest* 103:239–251

Panda DK, Miao D, Tremblay ML, Sirois J, Farookhi R, Henty GN, Goltzman D 2001 Targeted ablation of the 25-hydroxyvitamin D 1 α -hydroxylase enzyme: evidence for skeletal, reproductive, and immune dysfunction. *Proc Natl Acad Sci USA* 98:7498–7503

St-Arnaud R, Dardenne O, Prud'homme J, Hacking SA, Glorieux FH 2003 Conventional and tissue-specific inactivation of the 25-hydroxyvitamin D-1 α -hydroxylase (CYP27B1). *J Cell Biochem* 88:245–251

Lehste JR, Melsen F, Wellner M, Jansen P, Schlichting U, Renner-Muller I, Andreassen TT, Wolf E, Bachmann S, Nykjaer A, Willnow TE 2003 Hypocalcemia and osteopathy in mice with kidney-specific megalin gene defect. *FASEB J* 17:247–249 (Epub 2002 Dec 03)

Phosphatases

Suter A, Everts V, Boyde A, Jones SJ, Lullmann-Rauch R, Hartmann D, Hayman AR, Cox TM, Evans MJ, Meister T, von Figura K, Saftig P 2001 Overlapping functions of lysosomal acid phosphatase (LAP) and tartrate-resistant acid phosphatase (Acp5) revealed by doubly deficient mice. *Development* 128:4899–4910

Narisawa S, Frohlander N, Millan JL 1997 Inactivation of two mouse alkaline phosphatase genes and establishment of a model of infantile hypophosphatasia. *Dev Dyn* 208:432–446

Tesch W, Vandenbos T, Roschgr P, Fratzl-Zelman N, Klaushofer K, Beertsen W, Fratzl P 2003 Orientation of mineral crystallites and mineral density during skeletal

development in mice deficient in tissue nonspecific alkaline phosphatase. *J Bone Miner Res* 18:117–125

Cathepsin K

Saftig P, Hunziker E, Wehmeyer O, Jones S, Boyde A, Rommelskirch W, Moritz JD, Schu P, von Figura K 1998 Impaired osteoclastic bone resorption leads to osteopetrosis in cathepsin-K-deficient mice. *Proc Natl Acad Sci USA* 95:13453–13458

Gowen M, Lazner F, Dodds R, Kapadia R, Feild J, Tavariva M, Bertonecello I, Drake F, Zavorselk S, Tellis I, Hertzog P, Debouck C, Kola I 1999 Cathepsin K knockout mice develop osteopetrosis due to a deficit in matrix degradation but not demineralization. *J Bone Miner Res* 14:1654–1663

Prolactin

Heiman ML, Tinsley FC, Mattison JA, Hauck S, Bartke A 2003 Body composition of prolactin-, growth hormone, and thyrotropin-deficient Ames dwarf mice. *Endocrine* 20:149–154

Clement-Lacroix P, Ormandy C, Lepescheux L, Ammann P, Damotte D, Goffin V, Bouchard B, Amling M, Gaillard-Kelly M, Binart N, Baron R, Kelly PA 1999 Osteoblasts are a new target for prolactin: analysis of bone formation in prolactin receptor knockout mice. *Endocrinology* 140:96–105

Prostaglandin

Li X, Okada Y, Pilbeam CC, Lorenzo JA, Kennedy CR, Breyer RM, Raisz LG 2000 Knockout of the murine prostaglandin EP2 receptor impairs osteoclastogenesis *in vitro*. *Endocrinology* 141:2054–2061

Akhter MP, Cullen DM, Gong G, Recker RR 2001 Bone biomechanical properties in prostaglandin EP1 and EP2 knockout mice. *Bone* 29:121–125

Extra Cellular Matrix/Cartilage

Gowen LC, Petersen DN, Mansolf AL, Qi H, Stock JL, Tkalecic GT, Simmons HA, Crawford DT, Chidsey-Frink KL, Ke HZ, McNeish JD, Brown TA 2003 Targeted disruption of the osteoblast/osteocyte factor 45 gene (OF45) results in increased bone formation and bone mass. *J Biol Chem* 278:1998–2007 (Epub 2002 Nov 05)

Vu TH, Shipley JM, Bergers G, Berger JE, Helms JA, Hanahan D, Shapiro SD, Senior RM, Werb Z 1998 MMP-9/gelatinase B is a key regulator of growth plate angiogenesis and apoptosis of hypertrophic chondrocytes. *Cell* 93:411–422

Sahlman J, Inkinen R, Hirvonen T, Lammi MJ, Lammi PE, Nieminen J, Lapvetelainen T, Prockop DJ, Arita M, Li SW, Hyttinen MM, Helminen HJ, Puustjarvi K 2001 Premature vertebral endplate ossification and mild disc degeneration in mice after inactivation of one allele belonging to the Col2a1 gene for Type II collagen. *Spine* 26:2558–2565

Rosati R, Horan GS, Pinero GJ, Garofalo S, Keene DR, Horton WA, Vuorio E, de Crombrughe B, Behringer

RR 1994 Normal long bone growth and development in type X collagen-null mice. *Nat Genet* 8:129–135

Brunet LJ, McMahon JA, McMahon AP, Harland RM 1998 Noggin, cartilage morphogenesis, and joint formation in the mammalian skeleton. *Science* 280:1455–1457

Long F, Zhang XM, Karp S, Yang Y, McMahon AP 2001 Genetic manipulation of hedgehog signaling in the endochondral skeleton reveals a direct role in the regulation of chondrocyte proliferation. *Development* 128:5099–5108

Chellaiah M, Kizer N, Silva M, Alvarez U, Kwiatkowski D, Hruska KA 2000 Gelsolin deficiency blocks podosome assembly and produces increased bone mass and strength. *J Cell Biol* 148:665–678

Development

Orestes-Cardoso S, Nefussi JR, Lezot F, Oboeuf M, Pereira M, Mesbah M, Robert B, Berdal A 2002 Msx1 is a regulator of bone formation during development and postnatal growth: *in vivo* investigations in a transgenic mouse model. *Connect Tissue Res* 43:153–160

Satokata I, Ma L, Ohshima H, Bei M, Woo I, Nishizawa K, Maeda T, Takano Y, Uchiyama M, Heaney S, Peters H, Tang Z, Maxson R, Maas R 2000 Msx2 deficiency in mice causes pleiotropic defects in bone growth and ectodermal organ formation. *Nat Genet* 24:391–395

Sowa H, Kaji H, Canaff L, Hendy GN, Tsukamoto T, Yamaguchi T, Miyazono K, Sugimoto T, Chihara K 2003 Inactivation of menin, the product of the multiple endocrine neoplasia type 1 gene, inhibits the commitment of multipotential mesenchymal stem cells into the osteoblast lineage. *J Biol Chem* 278:21058–21069 (Epub 2003 Mar 20)

Kim IS, Otto F, Zabel B, Mundlos S 1999 Regulation of chondrocyte differentiation by Cbfa1. *Mech Dev* 80:159–170

Watanabe H, Yamada Y 1999 Mice lacking link protein develop dwarfism and craniofacial abnormalities. *Nat Genet* 21:225–229

Takagi T, Moribe H, Kondoh H, Higashi Y 1998 DeltaEF1, a zinc finger and homeodomain transcription factor, is required for skeleton patterning in multiple lineages. *Development* 125:21–31

Wang WF, Wang YG, Reginato AM, Plotkina S, Gridley T, Olsen BR 2002 Growth defect in Grg5 null mice is associated with reduced Ihh signaling in growth plates. *Dev Dyn* 224:79–89

Kato M, Patel MS, Levasseur R, Lobov I, Chang BH, Glass 2nd DA, Hartmann C, Li L, Hwang TH, Brayton CF, Lang RA, Karsenty G, Chan L 2002 Cbfa1-independent decrease in osteoblast proliferation, osteopenia, and persistent embryonic eye vascularization in mice deficient in Lrp5, a Wnt coreceptor. *J Cell Biol* 157:303–314

Miscellaneous

- Yoshida H, Hayashi S, Kunisada T, Ogawa M, Nishikawa S, Okamura H, Sudo T, Shultz LD, Nishikawa S 1990 The murine mutation osteopetrosis is in the coding region of the macrophage colony stimulating factor gene. *Nature* 345:442–444
- Zhang M, Xuan S, Bouxsein ML, von Stechow D, Akeno N, Faugere MC, Malluche H, Zhao G, Rosen CJ, Efstratiadis A, Clemens TL 2002 Osteoblast-specific knockout of the insulin-like growth factor (IGF) receptor gene reveals an essential role of IGF signaling in bone matrix mineralization. *J Biol Chem* 277:44005–44012 (Epub 2002 Sep 04)
- Bikle D, Majumdar S, Laib A, Powell-Braxton L, Rosen C, Beamer W, Nauman E, Leary C, Halloran B 2001 The skeletal structure of insulin-like growth factor I-deficient mice. *J Bone Miner Res* 16:2320–2329
- Gupta A, Tenenhouse HS, Hoag HM, Wang D, Khadeer MA, Namba N, Feng X, Hruska KA 2001 Identification of the type II Na(+)-Pi cotransporter (Npt2) in the osteoclast and the skeletal phenotype of Npt2^{-/-} mice. *Bone* 2001 29:467–476
- Garner SC, Pi M, Tu Q, Quarles LD 2001 Rickets in cation-sensing receptor-deficient mice: an unexpected skeletal phenotype. *Endocrinology* 142:3996–4005
- Wang Y, Spatz MK, Kannan K, Hayk H, Avivi A, Gorivodsky M, Pines M, Yayon A, Lonai P, Givol D 1999 A mouse model for achondroplasia produced by targeting fibroblast growth factor receptor 3. *Proc Natl Acad Sci USA* 96:4455–4460
- Silva IV, Cebotaru V, Wang H, Wang XT, Wang SS, Guo G, Devuyst O, Thakker RV, Guggino WB, Guggino SE 2003 The CIC-5 knockout mouse model of Dent's disease has renal hypercalciuria and increased bone turnover. *J Bone Miner Res* 18:615–623
- Sun L, Iqbal J, Dolgilevich S, Yuen T, Wu XB, Moonga BS, Adebajo OA, Bevis PJ, Lund F, Huang CL, Blair HC, Abe E, Zaidi M 2003 Disordered osteoclast formation and function in a CD38 (ADP-ribosyl cyclase)-deficient mouse establishes an essential role for CD38 in bone resorption. *FASEB J* 17:369–375
- Ohyama K, Chung CH, Chen E, Gibson CW, Misof K, Fratzl P, Shapiro IM 1997 p53 influences mice skeletal development. *J Craniofac Genet Dev Biol* 17:161–171
- Ho C, Conner DA, Pollak MR, Ladd DJ, Kifor O, Warren HB, Brown EM, Seidman JG, Seidman CE 1995 A mouse model of human familial hypocalciuric hypercalcemia and neonatal severe hyperparathyroidism. *Nat Genet* 11:389–394